### MADE EASY&NEXT IAS GROUP

PRESENT



Corporate Office: 44-A/1, Kalu Sarai, New Delhi 110016 | Web: www.meniit.com

Maximum Marks: 720 Time: 3 Hours



# **NEET (UG) - 2016**

### **IMPORTANT INSTRUCTIONS**

- 1. The Answer Sheet is inside the Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **side-1** and **side-2** carefully with blue/ black toll point pen only.
- 2. The test is of 3 hours duration and lest Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page / marking responses.
- 4. Rough work o to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must handover the Answer Sheet to the invigilator in the Room/Hall. The candidate are allowed to take away this Test Booklet with them.
- 6. The CODE forth it Booklet is **S**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Booklet. In case of discrepancy, the candidate should immediately report the matter to the invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your roll no. anywhere else except in the specified space in the Test Booklet /vAnswer Sheet.
- 8. Use of white fluid for correction is NOT permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admission Card to the Invigilator.
- 10. No candidates, without special permission of the Superintendent or Invigilator would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet the second time will be deemed not to have handed over Answer Sheet and dealt with as an unfair means ease.
- 12. Use of Electronic / Manual Calculator is prohibited
- 13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the Correct Test Booklet Code as given in the Test Booklet / Answer Sheet in the Attendance Sheet.

# **SECTION - I (BIOLOGY)**

**360 MARKS** 

- In a testcross involving  $F_1$  dihybrid flies, more parental-type offspring were produced than the recombinat-type offspring. This indicates:
  - (1) Both of the characters are controlled by more than one gene.
  - (2) The two genes are located on two different chromosomes
  - (3) Chromosomes failed to separate during meiosis.
  - (4) The two genes are linked and present on the same chromosome.
- **2.** Water soluble pigments found in plant cell vacuoles are:
  - (1) Anthocyanins
- (2) Xanthophylls
- (3) Chlorophylls
- (4) Carotenoids
- 3. Which of the following pairs of hormones are not antagonistic (having opposite effects) to each other?
  - (1) Relaxin Inhibin

(2) Parathormone - Calcitonin

(3) Insulin - Glucagon

- (4) Aldosterone Atrial Natriuretic factor
- **4.** Mitochondria and chloroplast are:
  - (a) semi-autonomous organelles.
  - (b) formed by division of pre existing organelles and they contain DNA but lack protein synthesizing machinery.

Which one of the following options is correct?

(1) Both (a) and (b) are false.

(2) Both (a) and (b) are correct.

(3) (b) is true but (a) is false.

- (4) (a) is true but (b) is false.
- 5. Which of the following is not a feature of the plasmids?
  - (1) Single stranded

(2) Independent replication

(3) Circular structure

- (4) Transferable
- A plant in your garden avoids photorespiratory losses, has improved water use efficiency, shows high rates of photosynthesis at high temperatures and has improved efficiency of nitrogen utilization. In which of the following physiological groups

would you assign this plant?

- (1) Nitrogen fixer
- $(2) C_3$
- (3) C<sub>4</sub>
- (4) CAM
- 7. Emerson's enhancement effect and Red drop have been instrumental in the discovery of:

Tip of nose

(1) Oxidative phosphorylation

(4) Transitional epithelium

- (2) Photophosphorylation and non-cyclic electron transport
- (3) Two photosystems opera ting simultaneously
- (4) Photophosphorylation and cyclic electron transport
- **8.** Which type of tissue correctly matches with its location?

Tissue Location
(1) Cuboidal epithelium Lining of stomach
(2) Smooth muscle Wall of intestine
(3) Areolar tissue Tendons

9.	When does the growth rate of a population following the logistic model equal zero? The logistic model is given as $dN/dt = rN(l-N/K)$ :						
		te is greater than birth rate.		(2) when N/K is exactly one.			
	(3) When N nears the carrying capacity of the habi		e habitat	(4) when N/K equals zero.			
10.	Which one of the following statements is not true?						
	(1) Stored pollen in liquid nitrogen can be used in the crop breeding programmes						
	(2) Tapetum helps in the dehiscence of anther						
	(3) Exine of pollen grains is made up of sporopollenin						
	(4) Pollen grains	(4) Pollen grains of many species cause severe allergies					
11.	Which one of the	following statements is wro	ong?				
	(1) Phycomycetes	are also called algal fungi.	(2) Cyanobacteria	are also called blue-green algae.			
	(3) Golden algae	are also called desmids.	(4) Eubacteria are	also called false bacteria.			
12.	The A vena curva	ture is used for bioassay of:					
	(1) Ethylene	<b>(2)</b> ABA	(3) GA <sub>3</sub>	<b>(4)</b> IAA			
13.	Which of the follo	owing structures is homolog	gus to the wing of a bin	rd?			
	(1) Flipper of Whale		(2) Dorsal fin of a	(2) Dorsal fin of a Shark			
	(3) Wing of a Mo	th	(4) Hind limb of F	(4) Hind limb of Rabbit			
14.	Blood pressure in	Blood pressure in the pulmonary artery is:					
	(1) less than that in the venae cavae. (2) same as that in the aorta.						
	(3) more than that	t in the carotid.	(4) more than that	in the pulmonary vein.			
15.	Fertilization in humans is practically feasible only if:						
	(1) the sperms are transported into cervix within 48 hrs of release of ovum in uterus.						
	(2) the sperms are transported into vagina just after the release of ovum in fallopian tube,						
	(3) the ovum and sperms are transported simultaneously to ampullary - isthmic junction of the fallopian tube.						
	(4) the ovum and	sperms are transported simu	ıltaneously to ampulla	ry - isthmic junction of the cervix.			
16.	In meiosis crossin	ng over is initiated at:	, )				
	(1) Diplotene	(2) Pachytene	(3) Leptotene	(4) Zygotene			
17.	Chrysophytes, Eu	glenoids, Dinoflagellates an	nd Slime moulds are in	ncluded in the kingdom:			
	(1) Animalia	(2) Monera	(3) Protista	<b>(4)</b> Fungi			
18.		n between successive stimul					
	(1) Tonus	(2) Spasm	(3) Fatigue	(4) Tetanus			
19.	<del>-</del>	ct statement on 'inhibin':	1.11.11.11.11.11.11.11.11.11.11.11.11.1	Y Y Y			
		(1) Is produced by nurse cells in testes and inhibits the secretion of LH.					
	` '	cretion of LH, FSH and Pro y granulose cells in ovary an		noffCII			
	•	y granulose cells in ovary an					
	(4) is produced by	g Grandrose cens in ovary an	ia minoris ine secretio	n or Em.			

20. Name the chronic respiratory disorder caused mainly by cigarette smoking:			ng:				
	(1) Respiratory alkalosis (2) Emphysema	(3) Asthma	(4) Respirator)' acidosis				
21.	Which of the following most appropriately descr	ibes haemophilia?					
	(1) Dominant gene disorder	(2) Recessive gene d iso	order				
	(3) X-linked recessive gene disorder	(4) Chromosomal disor	der				
22.	Select the correct statement:						
	(1) The leaves of gymnosperms are not well adapted to the second of the	oted to extremes of clim	ate				
	(2) Gymnosperms are both homosporous and het	(2) Gymnosperms are both homosporous and heterosporous					
	(3) Salvinia, Ginkgo and Pinuf all are gymnospe	rms					
	(4) Sequoia is one of the tallest trees						
23.	Which of the following is required as inducer(s)	for the expression of La					
	(1) lactose and galactose (2) glucose	(3) galactose	(4) lactose				
24.	A tall true breeding garden pea plant is crossed v F <sub>1</sub> plants were selfed the resulting genotypes wer		g garden pea plant. When the				
	(1) 3:1:: Dwarf: Tall						
	(2) 1:2:1:: Tall homozygous: Tall heterozygous : Dwarf						
	(3) 1:2:1:: Tall heterozygous: Tall homozygous : Dwarf						
	(4) 3:1:: Tall: Dwarf	(4) 3:1:: Tall: Dwarf					
25.	Which part of the tobacco plant is infected by M	eloidogyite incognita?	O'				
	<b>(1)</b> Root <b>(2)</b> Flower	(3) Leaf	(4) Stem				
26.	Which of the following is not a characteristic feature during mitosis in somatic cells?						
	(1) Synapsis	(2) Spindle fibres					
	(3) Disappearance of nucleolus	(4) Chromosome move	ment				
27.	Which of the following statements is not true for cancer cells in relation to mutations?						
	(1) Mutations inhibit production of telomerase.	(1) Mutations inhibit production of telomerase. (2) Mutations in proto-oncogenes accelerate the cell cycle					
	(3) Mutations destroy telomerase inhibitor.						
28.	One of the major components of cell wall of mos	st fungi is:					
	(1) Hemicellulose (2) Chitin	(3) Peptidoglycan	(4) Cellulose				
29.	Cotyledon of maize grain is called:	., .					
	(1) scutellum (2) plumule	(3) coleorhiza	(4) coleoptile				
30.	Which of the following would appear as the pior	neer organisms on bare r	ocks?				
	(1) Green algae (2) Lichens	(3) Liverworts	(4) Mosses				
31.	Changes in GnRH pulse frequency in females is	` '	• *				
	(1) progesterone and inhibin	(2) estrogen and proges	-				
	(3) estrogen and inhibin	(4) progesterone only					
	· · · · · · · · · · · · · · · · · · ·	, i 5					

<b>32.</b>	Antivenom injection contains preformed antibodies while polio drops that are administe	red into 1	the body
	contain:		

(1) Attenuated pathogens

(2) Activated pathogens

(3) Harvested antibodies

(4) Gamma globulin

- 33. Photosensitive compound in human eye is made up of:
  - (1) Transducin and Retinene(2) Guanosine and Retinol
  - (3) Opsin and Retinal (4) Opsin and Retinol
- 34. Specialised epidermal cells surrounding the guard cells are called:

(1) Lenticels

(2) Complementary cells

(3) Subsidiary cells

(4) Bulliform cells

- **35.** Which of the following features is not present in the Phylum - Arthropoda?
  - (1) Jointed appendages

(2) Chitinous exoskeleton

(3) Metameric segmentation

(4) Parapodia

- **36.** Reduction in pH of blood will:
  - (1) release bicarbonate ions by the liver.
- (2) reduce the rate of heart beat.
- (3) reduce the blood supply to the brain
- (4) decrease the affinity of hemoglobin with oxygen.
- 37. Which of the following characteristic features always holds true for the corresponding group of animals?

(1)	3 - chambered heart with one incompletely divided ventricle	Reptilia
(2)	Cartilaginous endoskeleton	Chondrichthyes
(3)	Viviparous	Mammalia
(4)	Possess a mouth with an upper and a lower jaw	Chordata

38. Match the terms in Column I with their description in Column II and choose the correct option:

Column I	Column II
(a) Dominance	(i) Many genes govern a single character
(b) Codominance	(ii) In a heterozygous organism only one allele expresses itself
(c) Pleiotropy	(iii) In a heterozygous organism both alleles express themselves fully
(d) Polygenic inheritance	(iv) A single gene influences many characters C

### Code:

	(a)	(b)	(c)	(d)
(1)	(iv)	(iii)	(i)	(ii)
(2)	(ii)	(i)	(iv)	(iii)
(3)	(ii)	(iii)	(iv)	(i)
(4)	(iv)	(i)	(ii)	(iii)

39.	A typical fat molecule	e is made up of:					
	(1) Three glycerol and three fatty add molecules						
	(2) Three glycerol molecules and one fatty acid molecule.						
	(3) One glycerol and t	hree fatty acid molecule	s				
	(4) One glycerol and o	one fatty acid molecule					
40.	Proximal end of the fi	lament of stamens attach	ned to the:				
	(1) Thalamus or petal	(2) Anther	(3) Connective	(4) Placenta			
41.	Which one of the follo	owing statements is wron	ng?				
	(1) Glycine is a sulph	(1) Glycine is a sulphur containing amino acid.					
	(2) Sucrose is a disacc	(2) Sucrose is a disaccharide.					
	(3) Cellulose is a poly	saccharide.					
	(4) Uracil is a pyrimic	line.					
42.	Water vapour comes out from the plant leaf through the stomatal opening. Through the same stomatal opening carbon dioxide diffuses into the plant during photosynthesis. Reason out the above statements using one of following options:						
	(1) One process occurs during day time, and the other at night.						
	(2) Both processes cannot happen simultaneously.						
	(3) Both processes can happen together because the diffusion coefficient of water and CO <sub>2</sub> is different						
	(4) The above processes happen only during nighttime.						
<b>43.</b>	A complex of ribosomes attached to a single strand of RNA is known as:						
	(1) Okazaki fragment	(2) Polysome	(3) Polymer	(4) Polypeptide			
<b>44.</b>	Which one of the following is a characteristic feature of cropland ecosystem?						
	(1) Ecological success	sion	(2) Absence of soil org	anisms			
	(3) Least genetic dive	rsity	(4) Absence of weeds				
<b>45.</b>	Which of the following is the most important cause of animals and plants being driven to extinction?						
	(1) Co - extinctions		(2) Over-exploitation				
	(3) Alien species inva	sion	(4) Habitat loss and fra	gmentation			
46.	In a chloroplast the hi	ghest number of protons	is found in:				
	(1) Antennae complex		(2) Stroma				
	(3) Lumen of thyiakoids		(4) Inter membrane space				
<b>47.</b>			of the techniques of DNA	A fingerprinting available at present?			
	(1) DNA - DNA hybr		(2) Polymerase chain reaction				
	(3) Zinc finger analys		(4) Restriction enzyme				
48.	The primitive prokary include the:	otes responsible for the	production of biogas from	the dung of ruminant animals,			
	(1) Eubacteria	(2) Halophiles	(3) Thermoaddophiles	(4) Methanogens			

- **49.** Which of the following features is not present in *Penplaneta americana*?
  - (1) Metamerically segmented body
  - (2) Schizocoelom as body cavity^
  - (3) Indeterminate and radial cleavage during embryonic development
  - (4) Exoskeleton composed of N-acetylglucosamine
- **50.** A system of rotating crops with legume or grass pasture to improve soil structure and fertility is called:
  - (1) Shifting agriculture (2) Ley farming
- (3) Contour farming
- (4) Strip farming
- **51.** Which of the following is wrongly matched in the given table?

	Microbe	Product	Application
(1)	Clostridium butylicum	Lipase	removal of oil stains
(2)	Trichoderma polysporum	Cyclosporin A	Immunosuppressive drug
(3)	Monascus purpureus	Statins	lowering of blood cholesterol
(4)	Streptococcus	Streptokinase	removal of clot from blood vessel

- **52.** In mammals, which blood vessel would normally carry largest amount of urea?
  - (1) Hepatic Portal Vein (2) Renal Vein
- (3) Dorsal Aorta
- (4) Hepatic Vein

- 53. Pick out the correct statements:
  - (a) Haemophilia is a sex-linked recessive disease
  - (b) Down's syndrome is due to aneuploidy.
  - (c) Phenylketonuria is an autosomal recessive gene disorder.
  - (d) Sickle cell anaemia is an X linked recessive gene disorder.
  - **(1)** (a), (b) and (c) are correct.

(2) (a) and (d) are correct.

(3) (b) and (d) are correct

- (4) (a), (c) and (d) are correct
- **54.** Which of the following guards the opening of hepatopancreatic duct into the duodenum?
  - (1) Sphincter of Oddi (
- (2) Semilunar valve
- (3) Deocaecal valve
- (4) Pyloric sphincter

- **55.** Microtubules are the constituents of:
  - (1) Centrosome, Nucleosome and Centrioles
- (2) Cilia, Flagella and Peroxisomes
- (3) Spindle fibres, Centrioles and Cilia
- (4) Centrioles, Spindle fibres and Chromatin
- **56.** The coconut water from tender coconut represents:
  - (1) Free nuclear endosperm
- (2) Endocarp

(3) Fleshy mesocarp

- (4) Free nuclear pro embryo
- 57. Tricarpellary, syncarpous gynoecium is found in flowers of:
  - (1) Poaceae
- (2) Liliaceae
- (3) Solanaceae
- (4) Fabaceae

- **58.** Which of the following is not a stem modification?
  - (1) Flattened structures of Opuntia
- (2) Pitcher of Nepenthes

(3) Thorns of citrus

(4) Tendrils of cucumber

59.	The taq polymerase enzyme is obtained from:						
	(1) Pseudomonas putida		(2) Thermtis aqttaticu	(2) Thermtis aqttaticus			
	(3) Thiobacillus ferroxidans		(4) Bacillus subtilis				
60.	Stems modified in	nto flat green organs perfo	orming the functions of leave	es are known as:			
	(1) Scales	(2) Cladodes	(3) Phyllodes	(4) Phylloclades			
61.	· ·	ates, the immune system city and it attacks self-cells	· ·	non-self. If this property is lost due to			
	(1) Active immur	nity	(2) Allergic response				
	(3) Graft rejection	n	(4) Auto-immune dise	rase			
<b>62.</b>	Nomenclature is	governed by certain unive	ersal rules.				
	Which one of the	following is contrary to t	he rules of nomenclature?				
	(1) When written	by hand, the names are to	be underlined				
	(2) Biological na	mes can be written in any	language				
	(3) The first word	l in a biological name repr	resents the genus name, and	the second is a specific epithet			
	(4) The names are	(4) The names are written in Latin and are italicised					
63.	In bryophytes and	d pteridophytes, transport	of male gametes requires:				
	(1) Water	<b>(2)</b> Wind	(3) Insects	(4) Birds			
64.	In context of Amniocentesis/ which of the following statement is incorrect?						
	(1) It can be used for detection of Cleft palate.						
	(2) It is usually done when a woman is between 14-16 weeks pregnant						
	(3) It is used for prenatal sex determination.						
	(4) It can be used	(4) It can be used for detection of Down syndrome.					
<b>65.</b>	In the stomach, gastric ad£ is secreted by the:						
	(1) acidic cells	(2) gastrin secretir	ng cells (3) parietal cells	(4) peptic cells			
66.	Spindle fibres att	ach on to:					
	(1) Kinetosome o	f the chromosome	(2) Telomere of the ch	romosome			
	(3) Kinetochore of the chromosome (4) Centromere of the chromosome						
<b>67.</b>	Which is the National Aquatic Animal of India?						
	(1) Sea - horse	(2) Gangetic shark	(3) River dolphin	(4) Blue whale			
68.	Which one of the	following cell organelles	is enclosed by a single mem	brane?			
	(1) Nuclei	(2) Mitochondria	(3) Chloroplasts	(4) Lysosomes			
69.	The two polypept	tides of human insulin are	linked together by:				
	(1) Disulphide br	idges (2) Hydrogen bond	ds (3) Phosphodiester bo	nd(4) Covalent bond			
70.		ollowing, all three are made	•				
	(1) Nitrogen, nicl	<u> </u>	(2) Boron, zinc, mang	anese			
	(3) Iron, copper,	· ·	(4) Molybdenum, mag				
	., , , , , , , , , , , , , , , , , , ,	-	. , ,	· · ·			

- 71. Which of the following statements is wrong for viroids?
  - (1) Their RNA is of high molecular weight
- (2) They lack a protein coat
- (3) They are smaller titan viruses
- (4) They cause infections
- **72.** Analogous structures are a result of:
  - (1) Stabilizing selection (2) Divergent evolution (3) Convergent evolution
- (4) Shared ancestry

- **73.** Select the incorrect statement:
  - (1) LH triggers secretion of androgens from the Leydig cells.
  - (2) FSH stimulates the Sertoli cells which help in sperrniogenesis.
  - (3) LH triggers ovulation in ovary
  - (4) LH and FSH decrease gradually during the follicular phase.
- 74. Which one of the following characteristics is not shared by birds and mammals?
  - (1) Warm blooded nature

(2) Ossified endoskeleton

(3) Breathing using lungs

(4) Viviparity

- **75.** Which of the following statements is not correct?
  - (1) Some reptiles have also been reported as pollinators in some plant species
  - (2) Pollen grains of many species can germinate on the stigma of a flower, but only one pollen tube of the same species grows into the style.
  - (3) Insects that consume pollen or nectar without bringing about pollination are called pollen/nectar robbers
  - (4) Pollen germination and pollen tube growth are regulated by chemical components of pollen interacting with those of the pistil.
- **76.** Seed formation without fertilization in flowering plants involves the process of:
  - (1) Apomixis
- (2) Sporulation
- (3) Budding
- (4) Somatic hybridization
- 77. Which of the following approaches does not give the defined action of contraceptive?

(1)	Vasectomy	prevents spermatogenesis	
(2)	Barrier methods	prevent fertilization	
(3)	Intra uterine devices	increase phagocytosis of sperms, suppress sperm motility and	
		fertilizing capacity of sperms	
(4)	Hormonal contraceptives	Prevent/retard entry of sperms, prevent ovulation and	
		fertilization	

- 78. The amino acid Tryptophan is the precursor for the synthesis of:
  - (1) Cortisol and Cortisone

- (2) Melatonin and Serotonin
- (3) Thyroxine and Triiodothyronine
- (4) Estrogen and Progesterone
- 79. A river with an inflow of domestic sewage rich in organic waste may result in:
  - (1) Death of fish due to lack of oxygen.
  - (2) Drying of the river very soon due to algal bloom.
  - (3) Increased population of aquatic food web organisms.
  - (4) An increased production of fish due to biodegradable nutrients.


80.	Gause's principle of competitive exclusion states that:					
	(1) Larger organisms exclude smaller ones through competition.					
	(2) More abundant species will exclude the less abundant species through competition.					
(3) Competition for the same resources excludes species having different food preferences.						
	(4) No two species can occupy the same niche, indefinitely for the same limiting resources.					
81.	Asthma may be attribut	ed to:				
	(1) accumulation of flu	id in the lungs	(2) bacterial infection of	f the lungs		
	(3) allergic reaction of	the mast cells in the lung	s (4) inflammation of the	trachea		
82.	The standard petal of a	papilionaceous corolla is	s also called:			
	(1) Corona	(2) Carina	(3) Pappus	(4) Vexfllum		
83.	Which of the following	is a restriction endonucl	ease?			
	(1) RNase	(2) Hind II	(3) Protease	(4) DNase		
84.	It is much easier for a s	mall animal to run uphill	I than for a large animal,	because:		
	(1) The efficiency of m	uscles in large animals is	s less than in the small ar	nimals.		
	(2) It is easier to carry a	, -		'.0'		
		ve a higher metabolic rat	e			
	` ′	a lower O <sub>2</sub> requirement				
85.	•	statements regarding the		O'		
	•		earth were non-green and	•		
	· ·		emoautotrophs that never	released oxygen.		
		s which one of the follow		(4) 1 (4)		
	(1) Both (a) and (b) are		(2) (a) is correct bu	· ·		
0.6	(3) (b) is correct but (a)		(4) Both (a) and (b) are			
86.	•			the field. He tells his teacher that this l plate and thus the cell is containing		
	more number of chrome		are is no formation of cer	i prace and thus the cen is containing		
		ling cells. This would res	sult in:			
	(1) Polyteny	(2) Aneuploidy	(3) Polyploidy	(4) Somaclonal variation		
87.	Depletion of which gas	in the atmosphere can le	ead to an increased incide	ence of skin cancers:		
	(1) Methane	(2) Nitrous oxide	(3) Ozone	(4) Ammonia		
88.	Joint Forest Manageme	nt Concept was introduc	ed in India during:			
	(1) 1990s	(2) 1960s	<b>(3)</b> 1970s	<b>(4)</b> 1980s		
89.	Which one of the follow	wing is the starter codon?	)			
	<b>(1)</b> UAG	<b>(2)</b> AUG	(3) UGA.	<b>(4)</b> UAA		
90.	The term ecosystem wa	as coined by:				
	(1) E. Wanning	<b>(2)</b> E.P.Odum	(3) A.G. Tansley	(4) E. Haeckel		

# **SECTION - II (PHYSICS)**

**180 MARKS** 

- **91.** What is the minimum velocity with which a body of mass m must enter a vertical loop of radius R so that it can complete the loop?
  - (1)  $\sqrt{5gR}$
- $(2) \sqrt{gR}$
- $(3) \sqrt{2gR}$
- **(4)**  $\sqrt{3gR}$
- **92.** If the magnitude of sum of two vectors is equal to the magnitude difference of the two vectors, the angle between these vectors is :
  - **(1)** 180°
- **(2)** 0°
- **(3)** 90°
- **(4)** 45°
- 93. At what height from the surface of earth the gravitation potential and the value of g are  $-5.4 \times 10^7$  J kg<sup>-2</sup> and 6.0ms<sup>-2</sup> respectively? Take the radius of earth as 6400 km:
  - (1) 2000 km
- **(2)** 2600 km
- (3) 1600 km
- (4) 1400 km
- 94. A long solenoid has 1000 turns. When a current of 4A flows through it, the magnetic flux linked with each turn of the solenoid is  $4 \times 10^{-3}$  Wb. The self-inductance of the solenoid is:
  - **(1)** 1 H
- **(2)** 4 H
- (3) 3 H
- (4) 2 H
- 95. An inductor 20 mH, a capacitor 50  $\mu$ F and a resistor 40  $\Omega$  are connected in series across a source of emf V = 10 sin 340 t. The power loss in A.C. circuit is:
  - **(1)** 0.89 W
- (2) 0.51 W
- (3) 0.67 W
- (4) 0.76 W
- 76. Two identical charged spheres suspended from a common point by two massless strings of lengths l, are initially at a distance d(d << l) apart because of their mutual repulsion. The charges begin to leak from both the spheres at a constant rate. As a result, the spheres approach each other with a velocity v. The v varies as a function of the distance x between the spheres, as:
  - (1)  $v \propto x^{-1}$
- **(2)**  $v \propto x^{\frac{1}{2}}$
- (3)  $v \propto x$
- **(4)**  $v \propto x^{-\frac{1}{2}}$

97.



A capacitor of  $2\mu F$  is charged as shown in the diagram. When the switch S is turned to position 2, the percentage of its stored energy dissipated is:

- **(1)** 80%
- **(2)** 0%
- **(3)** 20%
- **(4)** 75%
- 98. A particle moves so that its position vector is given by  $\vec{r} = \cos \omega t \ \hat{x} + \sin \omega t \ \hat{y}$ . Where  $\omega$  is a constant.


Which of the following is true?

- (1) Velocity is perpendicular to  $\vec{r}$  and acceleration is directed away from the origin.
- (2) Velocity and acceleration both are perpendicular to  $\vec{r}$ .
- (3) Velocity and acceleration both are parallel to  $\vec{r}$ .
- (4) Velocity is perpendicular to  $\vec{r}$  and acceleration is directed towards the origin.

- 99. From a disc of radius R and mass M, a circular hole of diameter R, whose rim passes through the centre is cut. What is the moment of inertia of the remaining part of the disc about a perpendicular axis, passing through the centre?
  - (1)  $9 \text{ MR}^2/32$
- (2)  $15 \text{ MR}^2/32$
- (3)  $13 \text{ MR}^2/32$
- (4)  $11 \text{ MR}^2/32$
- The ratio of escape velocity at earth  $(v_e)$  to the escape velocity at a planet  $(v_p)$  whose radius and mean density 100. are twice as that of earth is:
  - (1)  $1:\sqrt{2}$
- **(2)** 1:2
- (3)  $1:2\sqrt{2}$
- **(4)** 1:4
- 101. A potentiometer wire is 100 cm long and a constant potential difference is maintained across it. Two cells are connected in series first to support one another and then in opposite direction. The balance points are obtained at 50 cm and 10 cm from the positive end of the wire in the two cases. The ratio of emf's is:
  - (1) 3:2
- **(2)** 5:1
- **(3)** 5:4
- (4) 3:4
- 102. A siren emitting a sound of frequency 800 Hz moves away from an observer towards a cliff at a speed of 15 ms<sup>-1</sup>. Then, the frequency of sound that the observer hears in the echo reflected from the cliff is:

(Take velocity of sound in air =  $330 \text{ ms}^{-1}$ )

- (1) 885 Hz
- (2) 765 Hz
- (3) 800 Hz
- To get output 1 for the following circuit, the correct choice for the input is: 103.



(1) A = 1, B = 0, C = 1

(2) A = 0, B = 1, C = 0

(3) A = 1, B = 0, C = 0

- (4) A = 1, B = 1, C = 0
- 104. In a diffraction pattern due to a single slit of width 'a', the first minimum is observed at an angle 30° when light of wavelength 5000 Å is incident on the slit. The first secondary maximum is observed at an angle of:
- (2)  $\sin^{-1}\left(\frac{1}{4}\right)$  (3)  $\sin^{-1}\left(\frac{2}{3}\right)$  (4)  $\sin^{-1}\left(\frac{1}{2}\right)$
- 105. When a metallic surface is illuminated with radiation of wavelength  $\lambda$ , the stopping potential is V. If the same surface is illuminated with radiation of wavelength  $2\lambda$ , the stopping potential is  $\frac{V}{4}$ . The threshold wavelength for the metallic surface is:
  - (1)  $3\lambda$
- (2) 4λ
- **(3)** 5λ
- (4)  $\frac{5}{2}\lambda$
- When an  $\alpha$ -particle of mass 'm' moving with velocity 'v' bombards on a heavy nucleus of charge 'Ze', its 106. distance of closest approach from the nucleus depends on m as:
  - (1) m
- (2)  $\frac{1}{m}$
- (3)  $\frac{1}{\sqrt{m}}$
- (4)  $\frac{1}{m^2}$
- 107. Match the corresponding entries of column 1 with column 2. 'Where m is the magnification produced by the mirror]
  - Column 1
- Column 2
- (A) m = -2
- (a) Convex mirror

108.

109.

110.

111.

112.

113.

114.

(B)  $m = -\frac{1}{2}$ 

(C) m = +2

(b) Concave mirror

(c) Real image

(D) $m = +\frac{1}{2}$	(d) Virtual image		
(1) $A \rightarrow c$ and d;	$B \rightarrow b$ and d; $C \rightarrow b$ and	c; $D \rightarrow a$ and d	
(2) $A \rightarrow b$ and c;	$B \rightarrow b$ and c; $C \rightarrow b$ and	d; $D \rightarrow a$ and d	
(3) $A \rightarrow a$ and c;	$B \rightarrow a$ and d; $C \rightarrow a$ and b	$p$ ; $D \rightarrow c$ and $d$	
(4) $A \rightarrow c$ and d;	$B \rightarrow b$ and c; $C \rightarrow b$ and	d; $D \rightarrow b$ and c	
the magnitude of	_	tic energy of the particle	a constant tangential acceleration. What is a becomes equal to $8 \times 10^{-4}$ J by the end of
(1) $0.2 \text{ m/s}^2$	(2) $0.1 \text{ m/s}^2$	(3) $0.15 \text{ m/s}^2$	(4) $0.18 \text{ m/s}^2$
A small signal vo	Itage $V(t) = V_0 \sin \omega t$ is ap	plied across and ideal ca	apacitor C:
(1) Current I(t) le	eads voltage V(t) by 180°		
(2) Current I(t), 1	ags voltage V(t) by 90°		.0
(3) Over a full cy phase with vo	-	ot consume any energy	from the voltage source. Current I(t) is in
(4) Current I(t) is	in phase with voltage V(t)		·O'
A disk and a sphe	re of same radius but differ	ent masses roll off on ty	vo inclined planes of the same altitude and
length. Which one	e of the two objects gets to	the bottom of the plane	first ?
(1) Depends on the	heir masses	(2)	Disk
(3) Sphere		(4) Both reach at the	ne same time
			. Lengths of brass and steel rods are $l_{_{1}}$ and
			which one of the following relations holds
(1) $\alpha_1 l_1 = \alpha_2 l_2$	$(2)  \alpha_1 l_2 = \alpha_2 l_1$	(3) $\alpha_1 l_2^2 = \alpha_2 l_1^2$	(4) $\alpha_1^2 l_2 = \alpha_2^2 l_1$
A astronomical to		eyepiece of focal lengths	s 40 cm and 4 cm respectively. To view an
(1) 54.0 cm	(2) 37.3 cm	(3) 46.0 cm	<b>(4)</b> 50.0 cm
and passes throug		to a torque which produ	an axis which is perpendicular to its plane aces a constant angular acceleration of 2.0 ely.
<b>(1)</b> 3.0	<b>(2)</b> 8.0	<b>(3)</b> 7.0	<b>(4)</b> 6.0
A refrigerator wor	rks between 4°C and 30°C.	It is required to remove	600 calories of heat every second in order
to keep the tempe	rature of the refrigerated sp	ace constant. The power	r required is: (Take 1 cal = 4.2 Joules)
(1) 2365 W	<b>(2)</b> 2.365 W	(3) 23.65 W	<b>(4)</b> 236.5 W

- 115. A gas is compressed isothermally to half its initial volume. The same gas is compressed separately through an adiabatic process until its volume is again reduced to half. Then:
  - (1) Which of the case (whether compression through isothermal or through adiabatic process) requires more work will depend upon the atomicity of the gas.
  - (2) Compressing the gas isothermally will require more work to be done.
  - (3) Compressing the gas through adiabatic process will require more work to be done.
  - (4) Compressing the gas isothermally or adiabatically will require the same amount of work.
- The intensity at the maximum in a Young's double slit experiment is  $I_0$ . Distance between two slits is  $d = 5\lambda$ , 116. where  $\lambda$  is the wavelength of light used in the experiment. What will be the intensity in front of one of the slits on the screen place at a distance D = 10d?
- (2)  $I_0$  (3)  $\frac{I_0}{A}$
- Two non-mixing liquids of densities  $\rho$  and  $n\rho(n>1)$  are put in a container. The height of each liquid is h. A 117. solid cylinder of length L and density d is put in this container. The cylinder floats with its axis vertical and length pL(p < 1) in the denser liquid. The density d is equal to:

- (1)  $\{1+(n-1)\}p/\rho$  (2)  $\{1+(n+1)\}p/\rho$  (3)  $\{2+(n+1)\}p/\rho$  (4)  $\{2+(n-1)\}p/\rho$
- Consider the junction diode as ideal. The value of current flowing through AB is: 118.
  - (1)  $10^{-3}$  A
- (2) 0 A
- (3)  $10^{-2}$  A
- (4)  $10^{-1}$  A
- A car is negotiating a curved road of radius R. The road is banked at an angle  $\theta$ . The coefficient of friction 119. between the tyres of the car and the road is  $\mu_s$ . The maximum safe velocity on this road is:
  - (1)  $\sqrt{\frac{g}{R^2} \frac{\mu_s + \tan \theta}{1 \mu_s \tan \theta}}$  (2)  $\sqrt{gR^2 \frac{\mu_s + \tan \theta}{1 \mu_s \tan \theta}}$  (3)  $\sqrt{gR \frac{\mu_s + \tan \theta}{1 \mu_s \tan \theta}}$  (4)  $\sqrt{\frac{g}{R} \frac{\mu_s + \tan \theta}{1 \mu_s \tan \theta}}$

- A long straight wire of radius a carries a steady current I. The current uniformly distributed over its cross 120. section. The ratio of the magnetic fields B and B', at radial distances  $\frac{a}{2}$  and 2a respectively, form the axis of the wire is:
  - (1) 4
- (3)  $\frac{1}{2}$
- **(4)** 1
- Given the value of Rydberg constant is 10<sup>7</sup> m<sup>-1</sup>, the wave number of the last line of the Balmer series in 121. hydrogen spectrum will be:
  - (1)  $2.5 \times 10^7 \,\mathrm{m}^{-1}$
- (2)  $0.025 \times 10^4 \text{ m}^{-1}$
- (3)  $0.5 \times 10^7 \,\mathrm{m}^{-1}$
- (4)  $0.25 \times 10^7 \text{ m}^{-1}$
- Ii the velocity of a particle is  $v = At + Bt^2$ , where A and B are constants, then the distance travelled by it 122. between Is and 2s is:

- (1)  $\frac{A}{B} + \frac{B}{3}$  (2)  $\frac{3}{2}A + 4B$  (3) 3A + 7B (4)  $\frac{3}{2}A + \frac{7}{3}B$
- 123. The angle of incidence for a rayof lightata refracting surface of a prism is 45°. The angle of prism is 60°. If the ray suffers minimum deviation through the prism, the angle of minimum deviation and refractive index of the material of the prism respectively, are

<i>(</i> 1)	30°;	1
(1)	50,	$\sqrt{2}$

(2) 
$$45^{\circ}; \frac{1}{\sqrt{2}}$$

**(3)** 
$$30^{\circ}; \sqrt{2}$$

**(4)** 
$$45^{\circ}; \sqrt{2}$$

124. The molecules of a given mass of a gas have r.m.s. velocity of 200 ms<sup>-1</sup> at  $27^{\circ}$ C and  $1.0 \times 10^{5}$  Nm<sup>-2</sup> 124. pressure. When the temperature and pressure of the gas are respectively,  $127^{\circ}$ C and  $0.05 \times 10^{5}$  Nm<sup>-2</sup>. the r.m.s. velocity of its molecules in ms<sup>-1</sup> is:

(1) 
$$\frac{100}{3}$$

**(2)**  $100\sqrt{2}$ 

(3)  $\frac{400}{\sqrt{3}}$ 

(4)  $\frac{100\sqrt{2}}{3}$ 

An air column, closed at one end and open at the other, resonates with a tuning fork when the smallest length 125. of the column is 50 cm. The next larger length of the column resonating with the same tuning fork is:

(1) 200 cm

(2) 66.7 cm

(3) 100 cm

(4) 150 cm

126. The magnetic susceptibility is negative for:

(1) paramagnetic and ferromagnetic materials

(2) diamagnetic material only

(3) paramagnetic material only

(4) ferromagnetic material only

127. An electron or mass m and a photon have same energy E The ratio of de-Broglie wavelengths associated with

(1) 
$$\frac{1}{c} \left(\frac{2m}{E}\right)^{\frac{1}{2}}$$
 (2)  $\frac{1}{c} \left(\frac{E}{2m}\right)^{\frac{1}{2}}$  (3)  $\left(\frac{E}{2m}\right)^{\frac{1}{2}}$ 

A body of mass 1 kg begins to move under the action of a time dependent force  $\vec{F} = (2t\hat{i} + 2t^2\hat{j})N$ , where  $\hat{i}$  and 128. i are unit vectors along x and y axis. What power will be developed by the force at the time t?

(1)  $(2t^7 + 3t^5)W$ 

- (2)  $(2t^2+3t^3)W$
- (3)  $(2t^2+4t^4)$  W (4)  $(2t^3+3t^4)$ W
- The charge flowing through a resistance R varies with time t as  $Q=at-bt^2$ , where a and b are positive 129. constants. The total heat produced in R is:

(1)  $\frac{a^3 R}{b}$ 


- (2)  $\frac{a^3 R}{6h}$  (3)  $\frac{a^3 R}{3h}$
- (4)  $\frac{a^3 R}{2 h}$
- A npn transistor is connected in common emitter configuration in a given amplifier. A load resistance of 800 130.  $\Omega$  is connected in the collector circuit and the voltage drop across-it is 0.8 V. If the current amplification factor is 0.96 and the input resistance

of the circuit is 192  $\Omega$ , the voltage gain and the power gain of the amplifier will respectively be:

- **(1)** 4,3.69
- **(2)** 4, 3.84
- **(3)** 3.69,3.84
- **(4)** 4, 4
- A piece of ice falls from a height h so that it melts com pletely. Only one quarter of the heat produced is 131. absorbed by the ice and all energy of ice gets converted into heat during its fall. The value of h is: [Latent heat of ice is  $3.4 \times 10^5$  J/ kg and g = 10 N/ kg]

(1) 68 km

- **(2)** 34km
- (3) 544km
- (4) 136 km
- 132. A square loop A BCD carrying a current i, is placed near and coplanar with a long straight conductor XV carrying a current I, the net force on the loop will



- (1)  $\frac{\mu_0 \text{IiL}}{2\pi}$
- (2)  $\frac{2\mu_0 \text{Ii}}{2\pi}$
- (3)  $\frac{\mu_0 \text{Ii}}{2\pi}$
- (4)  $\frac{2\mu_0 \text{liL}}{3\pi}$
- A uniform rope of length L and mass m<sub>1</sub> hangs vertically from a rigid support. A block of mass m<sub>2</sub>, is attached 133. to the free end of the rope. A transverse pulse of wavelength  $\lambda_1$  is produced at the lower end of the rope. The wavelength of the puLse when it reaches the top of the rope is  $A_2$ . The ratio  $\lambda_2/\lambda_1$  is:
- (2)  $\sqrt{\frac{m_1}{m_2}}$  (3)  $\sqrt{\frac{m_1 + m_2}{m_2}}$  (4)  $\sqrt{\frac{m_2}{m_1}}$
- A black body is at a temperature of 5760 K. The energy of radiation emitted by the body at wavelength 250 134. nm is  $U_1$ , at wavelength 500 nm is  $U_2$  and that at 1000 nm is  $U_3$ . Wien's constant,  $b = 2.88 \times 10$  nmK. Which of the following is correct?
  - (1)  $U_2 > U_1$
- (2)  $U_1 = 0$
- (3)  $U_3 = 0$
- (4)  $U_1 > U_2$
- Out of the following options which one can be used to produces propagating electromagnetic wave? 135.
  - (1) An accelerating charge

(2) A charge moving at constant velocity

(3) A stationary charge


(4) A chargeless particle

# **SECTION - III (CHEMISTRY)**

**180 MARKS** 

- 136.<sup>E</sup> Which one of the following characteristics is associated with adsorption?
  - (1)  $\Delta G$  and  $\Delta S$  are negative but  $\Delta H$  is positive (2)  $\Delta G$  is negative but  $\Delta H$  and  $\Delta S$  are positive
  - (3)  $\Delta G$ ,  $\Delta H$  and  $\Delta S$  all are negative
- (4)  $\Delta G$  and  $\Delta H$  are negative but  $\Delta S$  is positive
- The pressure of H<sub>2</sub> required to make the potential of H<sub>2</sub>-electrode zero in pure water at 298 K is 137.<sup>M</sup>
  - (1)  $10^{-4}$  atm
- (2)  $10^{-14}$  atm
- (3)  $10^{-12}$  atm
- (4)  $10^{-10}$  atm
- 138.<sup>E</sup> The addition of a catalyst during a chemical reaction alters which of the following quantities?
- (1) Activation energy (2) Entropy
- (3) Internal energy
- (4) Enthalpy

- 139.<sup>E</sup> For the following reactions
  - (a)  $CH_3CH_2CH_2Br + KOH \rightarrow CH_3CH = CH_2 + KBr + H_2O$
  - (b)  $H_3C$   $CH_3$  + KOH + KBr OH



Which of the following statements is correct?

(1) (a) is substitution, (b) and (c) are addition reactions.

140.<sup>E</sup>

 $141.^{M}$ 

142.<sup>E</sup>

143.<sup>E</sup>

NEET 2010   Taper CODE . 1
(2) (a) and (b) are elimination reactions and (c) is addition reaction.
(3) (a) is elimination, (b) is substitution and (c) is addition reaction.
(4) (a) is elimination, (b) and (c) are substitution reactions.
The product formed by the reaction of an aldehyde with a primary amine is
(1) Aromatic acid (2) Schiff base (3) Ketone (4) Carboxylic acid
The correct statement regarding the basicity of arylamines is
(1) Arylamines are generally more basic than alkylamines, because the nitrogen atom in arylamines is sphybridized.
(2) Arylamines are generally less basic than alkylamines because the nitrogen lone-pair electrons are delocalized by interaction with the aromatic ring $\pi$ electron system.
(3) Arylamines are generally more basic than alkylamines because the nitrogen lone-pair electrons are not delocalized by interaction with the aromatic ring $\pi$ electron system.
(4) Arylamines are generally more basic than alkylamines because of aryl group.
Equal moles of hydrogen and oxygen gases are placed in a container with a pin-hole through which both can escape. What fraction of the oxygen escapes in the time required for one-half of the hydrogen to escape?
<b>(1)</b> 1/2 <b>(2)</b> 1/8 <b>(3)</b> 1/4 <b>(4)</b> 3/8
The correct statement regarding the comparison of staggered and eclipsed conformations of ethane, is
(1) The staggered conformation of ethane is more stable than eclipsed conformation, because staggered conformation has no torsional strain.
(2) The staggered conformation of ethane is less stable than eclipsed conformation, because staggered conformation has torsional strain.
(3) The eclipsed conformation of ethane is more stable than staggered conformation, because eclipsed conformation has no torsional strain.
(4) The eclipsed conformation of ethane is more stable than staggered conformation even through the eclipsed conformation has torsional strain.
In which of the following options the order of arrangement does not agree with the variation of property

144.<sup>M</sup> indicated against it?

(1) Li  $\leq$  Na  $\leq$  K  $\leq$  Rb (increasing metallic radius)

- (2)  $Al^{3+} < Mg^{2+} < Na^{+} < F^{-}$  (increasing ionic size)
- (3) B < C < N < O (increasing first ionization enthalpy)
- (4) I < Br < Cl < F (increasing electron gain enthalpy)

145.<sup>E</sup> The rate of a first-order reaction is  $0.04 \text{ mol } 1^{-1}\text{s}^{-1}$  at 10 seconds are  $0.03 \text{ mol } 1^{-1}\text{s}^{-1}$  at 20 seconds after initiation of the reaction. The half-life period of the reaction is

- (1) 54.1 sec
- **(2)** 24.1 sec
- **(3)** 34.1 sec
- **(4)** 44.1 sec

**146.** When copper is heated with conc. HNO<sub>3</sub> it produces:

(1)  $Cu(NO_3)_2$  and  $N_2O$ 

(2)  $Cu(NO_3)_2$  and  $NO_2$ 

(3)  $Cu(NO_3)_2$  and NO

(4)  $Cu(NO_3)_2$ , NO and  $NO_2$ 

147.<sup>E</sup> In a protein molecule, various amino acids are linked together by

- (1) dative bond (2)  $\alpha$ -glycosidic bond (3)  $\beta$ -glycosidic bond (4) peptide bond
- 148.<sup>E</sup> Fog is a colloidal solution of
  - (1) Gas in gas
- (2) Liquid in gas
- (3) Gas in liquid
- (4) solid in gas
- 149.<sup>M</sup> Match items of Column I with the items Column II and assign the correct code

### Column I

- (a) Cyanide process
- (b) Froth floatation process
- (c) Electrolytic reduction
- (d) Zone refining

#### Column II

- (i) Ultrapure Ge
- (ii) Dressing of ZnS
- (iii) Extraction of A
- (iv) Extraction of A
- (v) Purification of

### Code:

- (a) (b)
- (c)
- (v) (i)

(d)

(i)


(v)

- (1) (iii) **(2)** (iv)
- (iv) (ii)
- (iii)
- (3) (ii)
- (iii)
- (i)
- **(4)** (i)
- (ii)
- (iii) (iv)
- 150.<sup>M</sup> Which one given below is a non-reducing sugar?
  - (1) Sucrose
- (2) Maltose
- (3) Lactose
- (4) Glucose
- 151.<sup>E</sup> The correct statement regarding RNA and DNA respectively is:
  - (1) The sugar component in RNA 2'-deoxyribose and the sugar component DNA is arabinose.
  - (2) The sugar component in RNA is arabinose and the sugar component in DNA is 2'-deoxyribose
  - (3) The sugar component in RNA is ribose the sugar component in DNA 2'-deoxyribose.
  - (4) The sugar component in RNA is arabinose and the sugar component in DNA is ribose
- 152.<sup>E</sup> The correct thermodynamic conditions for spontaneous reaction at all temperatures is

  - (1)  $\Delta H < 0$  and  $\Delta S < 0$  (2)  $\Delta H <$  and  $\Delta S = 0$  (3)  $\Delta H > 0$  and  $\Delta S < 0$  (4)  $\Delta H < 0$  and  $\Delta S > 0$
- 153.<sup>D</sup> Which is the correct statement for the given acids?
  - (1) Phosphinic acid is a diprotic acid while phosphonic acid is a monoprotic acid
  - (2) Phosphinic acid is a monoprotic acid while phosphonic acid is a diprotic acid
  - (3) Both are diprotic acids

- (4) Both are triprotic acids.
- 154.<sup>M</sup> MY and NY<sub>3</sub>, two nearly insoluble salts, have the same  $K_{sp}$  values of  $6.2 \times 10^{-13}$  at room temperature. Which statement would be true in regard to MY and NY<sub>3</sub>?
  - (1) The addition of the salt of KY to solution of MY and NY<sub>3</sub> will have no effect on their solubilities.
  - (2) The molar solubilities of MY and NY<sub>3</sub> in water are identical.
  - (3) The molar solubility of MY in water is less than that of NY<sub>3</sub>.
  - (4) The salts MY and NY<sub>3</sub> are more soluble in 0.5 M KY than in pure water.
- 155.<sup>E</sup> Which of the following is an analgesic?

	(1) Chloromycetin	(2) Novalgin	(3)	Penicillin	(4) Streptomycin		
156. <sup>E</sup>	The pair of electrons in	the given carbanion, CI	$H_3C =$	■ C <sup>□</sup> is present in wh	nich of the following orbitals?		
	(1) sp	<b>(2)</b> 2p	(3)	$sp^3$	(4) $sp^2$		
157. <sup>M</sup>	Among the following,	the correct order of acidi	ty is				
	<b>(1)</b> HClO <sub>4</sub> < HClO <sub>2</sub> <	HClO < HClO <sub>3</sub>	(2)	$HClO_3 < HClO_4 < 1$	HClO <sub>2</sub> < HClO		
	(3) HClO < HClO <sub>2</sub> < H	$HClO_3 < HClO_4$	(4)	$HClO_2 < HClO < H$	$IClO_3 < HClO_4$		
158. <sup>E</sup>	Which one of the follow	wing statements is correct	et wh	en SO <sub>2</sub> is passed thre	ough acidified K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> solution?		
	(1) Green Cr <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> is formed.		(2) The solution turns blue				
	(3) The solution is decolorized			(4) SO <sub>2</sub> is reduced			
159. <sup>E</sup>	Predict the correct orde	er among the following					
	(1) lone pair – bond pa	air > bond pair – bond pa	ir > 1	lone pair – lone pair			
	(2) lone pair – lone pair	ir > lone pair – bond pair	· > bo	ond pair – bond pair			
	(3) lone pair – lone pair	ir > bond pair – bond pai	r > 10	one pair – bond pair			
	(4) bond pair – bond p	air > lone pair – bond pa	ir > ]	lone pair – lone pair			
160. <sup>E</sup>	Two electrons occupying the same orbital are distinguished by				`.O`		
	(1) Spin quantum number		(2) Principal quantum number				
	(3) Magnetic quantum	number	(4) Azimuthal quantum number				
161. <sup>M</sup>	The product obtained a	s a result of a reaction of	nitr	ogen with CaC2 is	O'		
	(1) Ca <sub>2</sub> CN	(2) Ca(CN) <sub>2</sub>	(3)	CaCN	(4) CaCN <sub>3</sub>		
162. <sup>D</sup>	Natural rubber has						
	(1) Random cis-and tra	ans-configuration	(2)	All cis-configuration	n		
	(3) All trans-configura	ation	(4)	Alternate cis-and tr	ans – configuration		



Can be classified as

- (1) Williamson alcohol synthesis reaction
- (2) Williamson ether synthesis reaction
- (3) Alcohol formation reaction
- **(4)** Dehydration reaction
- Lithium has a bcc structure. Its density is 530 kg m<sup>-3</sup> and its atomic mass is 6.94 g mol<sup>-1</sup>. Calculate the edge 165.<sup>M</sup> length of a unit cell of Lithium metal. ( $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ )

163. Which one of the following orders is correct for the bond dissociation enthalpy of halogen molecules? (1)  $F_2 > Cl_2 > Br_2 > I_2$  (2)  $I_2 > Br_2 > Cl_2 > F_2$  (3)  $Cl_2 > Br_2 > F_2 > I_2$  (4)  $Br_2 > I_2 > F_2 > Cl_2$ 

- (1) 264 pm
- **(2)** 154 pm
- (3) 352 pm
- (4) 527 pm


- The ionic radii of A<sup>+</sup> and B<sup>-</sup> ions are  $0.98 \times 10^{-10}$  m and  $1.81 \times 10^{-10}$  m. The coordination number of each ion 166.<sup>E</sup> in AB is
  - (1) 2
- **(2)** 6
- (3) 4
- **(4)** 8
- 167.<sup>E</sup> At 100°C the vapor pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If  $K_b = 0.52$ , the boiling point of this solution will be
  - (1) 103°C
- (2) 101°C
- (3) 100°C
- (4) 102°C
- 168.<sup>D</sup> The electronic configurations of Eu (Atomic No. 63), Gd (Atomic No. 64) and Tb (atomic No. 65) are
  - (1)  $[Xe]4f^76s^2$ ,  $[Xe]4f^75d^16s^2$  and  $[Xe]4f^96s^2$
  - (2)  $[Xe]4f^76s^2$ ,  $[Xe]4f^86d^16s^2$  and  $[Xe]4f^85d^16s^2$
  - (3)  $[Xe]4f^65d^16s^2$ ,  $[Xe]4f^75d^16s^2$  and  $[Xe]4f^96s^2$
  - (4)  $[Xe]4f^65d^16s^2$ ,  $[Xe]4f^75d^16s^2$  and  $[Xe]4f^85d^16s^2$
- 169.<sup>E</sup> Which of the following statements about hydrogen is incorrect?
  - (1) Dihydrogen does not act as a reducing agent
  - (2) Hydrogen has three isotopes of which tritium is the most common
  - (3) Hydrogen never acts as cation in ionic salts.
  - (4) Hydronium ion, H<sub>3</sub>O<sup>+</sup> exists freely in solution.
- In the reaction  $H C \equiv CH \xrightarrow{(1) \text{NaNH}_2/\text{liq.NH}_3} X \xrightarrow{(2) \text{CH}_3 \text{CH}_2 \text{Br}} Y$ , X and Y are 170.<sup>M</sup>
  - (1) X = 1-Butyne; Y = 2-Hexyne
- (2) X = 1-Butyne; Y = 3-Hexyne
- (3) X = 2-Butyne; Y = 3-Hexyne
- (4) X = 2-Butyne; Y = 2-Hexyne
- Consider the following liquid vapour equilibrium. 171.<sup>M</sup>


Liquid Vapour


Which of the following relations is correct?


- (1)  $\frac{d \ln P}{dT} = \frac{\Delta H_v}{RT^2}$

- (2)  $\frac{d \ln G}{dT^2} = \frac{\Delta H_v}{RT^2}$  (3)  $\frac{d \ln P}{dT} = \frac{-\Delta H_v}{RT}$  (4)  $\frac{d \ln P}{dT^2} = \frac{-\Delta H_v}{T^2}$
- $172.^{M}$ Which of the following statements about the composition of the vapour over an ideal 1: 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C. (Given, Vapour pressure data at 25°C, benzene = 12.8 kPa, toluene = 3.85 kPa)
  - (1) Not enough information is given to make a prediction
  - (2) The vapour will contain a higher percentage of benzene
  - (3) The vapour will contain a higher percentage of toluene
  - (4) The vapour will contain equal amounts of benzene and toluene.
- Which of the following biphenyls is optically active? 173.<sup>D</sup>









	(1) Aluminium isopropoxide (2	) Acetone
	(3) Ozone (4	$MnO_2$
175. <sup>D</sup>	The correct statement regarding a carbonyl compound	nd with a hydrogen atom on its alpha carbon, is
	(1) A carbonyl compound with a hydrogen atom or enol and this process is known as keto-enol tau	its alpha-carbon rapidly equilibrates with its corresponding tomerism.
	(2) A carbonyl compound with a hydrogen atom or enol.	n its alpha-carbon never equilibrates with its corresponding
		n its alpha-carbon rapidly equilibrates with its corresponding one equilibration.
	(4) A carbonyl compound with a hydrogen atom or enol and this process is known as carbonylation	its alpha-carbon rapidly equilibrates with its corresponding
176. <sup>E</sup>	Consider the molecules CH <sub>4</sub> , NH <sub>3</sub> and H <sub>2</sub> O. Which	of the given statements is false?
	(1) The $H - C - H$ bond angle in $CH_4$ is larger than	
	(2) The H $-$ C $-$ H bond angle in CH <sub>4</sub> ,	the H - N - H bond angle in NH <sub>3</sub> , and the
	$H - O - H$ bond angle in $H_2O$ are all greater that	
	(3) The $H - O - H$ bond angle in $H_2O$ is larger than	
4 == E	(4) The $H - O - H$ bond angle in $H_2O$ is smaller th	
177. <sup>E</sup>	Match the compounds given in Column I with the correct option	hybridization and shape given in Column II and mark the
	Column I Column II	
	(a) XeF <sub>6</sub> (i) Distorted octahedral	
	(b) XeO <sub>3</sub> (ii) Square planar (c) XeOF <sub>4</sub> (iii) Pyramidal	
	Code	
	(d) XeF <sub>4</sub> (iv) Square pyramidal  Code  (a) (b) (c) (d)  (1) (iv) (i) (ii) (iii)  (2) (i) (iii) (iv) (ii)  (3) (i) (ii) (iv) (iii)  (4) (iv) (iii) (i) (ii)	
	(1) (iv) (i) (ii) (iii) (2) (i) (iii) (iv) (ii)	
	(3) (i) (ii) (iv) (iii)	
150 M		
178. <sup>M</sup>	the mixture, the rate of nitration will be	. H <sub>2</sub> SO <sub>4</sub> and HNO <sub>3</sub> . If a large amount of KHSO <sub>4</sub> is added to
		Slower (4) Unchanged
179. <sup>M</sup>		(i) Chemangea
	(1) Mg <sup>2+</sup> ions are important in the green parts of pl	ants.
	(2) Mg <sup>2+</sup> ions form a complex with ATP (3	Ca <sup>2+</sup> ions are important in blood clotting
	(4) Ca <sup>2+</sup> ions are not important in maintaining the r	egular beating of the heart.
180. <sup>D</sup>		
	(1) $\left[ \text{Mn(CO)}_{6} \right]^{+}$ (2) $\text{Ni(CO)}_{4}$ (3)	$\left[\operatorname{Co(CO)}_{\cdot}\right]^{\ominus} \qquad \qquad \textbf{(4)} \left[\operatorname{Fe(CO)}_{\cdot}\right]^{2^{-}}$
	( ) [ ( )6] ( ) ( )4	/ L ( /4]

174.<sup>M</sup> Which of the following reagents would distinguish cis-cyclopenta-1, 2-diol from the trans-isomer?

**NEET 2016 : Paper Code** 



# **ANSWER KEY**

BIOLOGY			PHYSICS		CHEMISTRY		
1.	(4)	46.	(3)	91.	(1)	136.	(3)
2.	(1)	47.	(3)	92.	(3)	137.	(2)
3.	(1)	48.	(4)	93.	(2)	138.	(1)
4.	(4)	49.	(3)	94.	(1)	139.	(3)
5.	(1)	50.	(2)	95.	(2)	140.	(2)
6.	(3)	51.	(1)	96.	(4)	141.	(2)
7.	(3)	52.	(4)	97.	(1)	142.	(2)
8.	(2)	53.	(1)	98.	(4)	143.	(1)
9.	(2)	54.	(1)	99.	(3)	144.	(3 & 4)
10.	(2)	55.	(3)	100.	(3)	145.	(2)
11.	(4)	56.	(1)	101.	(1)	146.	(2)
12.	(4)	57.	(2)	102.	(4)	147.	(4)
13.	(1)	58.	(2)	103.	(1)	148.	(2)
14.	(4)	59.	(2)	104.	(1)	149.	(2)
15.	(3)	60.	(4)	105.	(1)	150.	(1)
16.	(2)	61.	(4)	106.	(2)	151.	(3)
17.	(3)	62.	(2)	107.	(2)	152.	(4)
18.	(4)	63.	(1)	108.	(2)	153.	(2)
19.	(3)	64.	(1)	109.	(3)	154.	(3)
20.	(2)	65.	(3)	110.	(3)	155.	(2)
21.	(3)	66.	(3)	111.	(1)	156.	(1)
22.	(4)	67.	(3)	112.	(1)	157.	(3)
23.	(4)	68.	(4)	113.	(2)	158.	(1)
24.	(2)	69.	(1)	114.	(4)	159.	(2)
25.	(1)		Incorrect	115.	(3)	160.	(1)
26.	(1)		Question	116.	(1)	161.	(Bonus)
27.	(1)		(Bonus)	117.	(1)	162.	(2)
28.	(2)	71.	(1)	118.	(3)	163.	(3)
29.	(1)	72.	(3)	119.	(3)	164.	(2)
30.	(2)	73.	(4)	120.	(4)	165.	(3)
31.	(2)	74.	(4)	121.	(4)	166.	(2)
32.	(1)	75.	(2)	122.	(4)	167.	(2)
33.	(3)	76.	(1)	123.	(3)	168.	(1)
34. 35.	(3)	77.	(1)	124. 125	(3)	169.	(1 & 2)
35. 36.	(4)	78.	(2)	125.	(4)	170.	(2)
30. 37.	(4) (2)	79. 80.	(1)	126. 127.	(2)	171. 172.	(1)
			(4)		(2) (1)		(2)
38. 39.	(3) (3)	81. 82.	(3)	128. 129.	(1) (2)	173. 174.	(3)
39. 40.	(1)	83.	(4) (2)	130.	(2) (2)	174. 175.	(2) (1)
40.	(1)	84.	(3)	130. 131.	(4)	175. 176.	(3)
41.	(3)	85.	(4)	131. 132.	(2)	170.	(2)
43.	(2)	86.	(3)	133.	(3)	177.	(3)
44.	(3)	87.	(3)	134.	(4)	178.	(4)
45.	(4)	88.	(4)	135.	(1)	180.	(4)
	(.)	89.	(2)	133.	(*)	100.	(')
		90.	(3)				
L		1	V /				

# **HINTS & SOLUTION**

### **BIOLOGY**

1. (4)

**Sol.:** Linkage increase incidence of parentals.

Hint: Chapter Principles of Inheritance and Variation

NCERT page no.83

2. (1)

Anthocyanins are water soluble vacuolar pigments

Hint: Chapter Photosynthesis in higher plants

Not given in NCERT

3. (1)

**Sol.:** Relaxin hormone causes relaxation of pelvic ligaments during parturition were as inhibin has negative

feedback effect with FSH

Chapter: Chemical co-ordination and integration

OUT of NCERT

4. (4)

**Sol.:** Both are semi autonomous and have protein synthesizing machinery including ribosomes.

Hint: Chapter cell the unit of life

NCERT page no.134

**5.** (1)

**Sol.:** Plasmids are ds-DNA molecules that are separate from a cells nucleoid in prokaryotes.

Chapter: Biotechnology Principles and Processes

Plasmid is double stranded DNA, Autonomously replicating unit, found in prokaryotes.

**6.** (3)

**Sol.:** Reduction of photorespiration in C<sub>4</sub> plants enhances both yield and nitrogen efficiency.

Hint: Chapter Photosynthesis in higher plants

NCERT page no.90

7. (3)

Hint: Chapter Photosynthesis in higher plants

Not given in NCERT

8. (2)

**Sol.:** Smooth muscles are involuntary muscles present in wall of intestine.

Chapter: Structural organization in Animals

NCERT Page 101- last para, 103- 1<sup>st</sup> and 2<sup>nd</sup> para, 104- 2<sup>nd</sup> para

9. (2)

Hint: Chapter Organisms and Population

NCERT page no.231

10. (2)

**Sol.:** Tapetum is nutritive layer of anther.

Hint: Chapter Sexual Reproduction in flowering plants

NCERT page no.22 to 24

11. (4)

**Sol.**; Eubacteria literally means true bacteria.

Hint: Chapter Biological classification

NCERT page no.19

**12.** (4)

**Sol.:** Avena curvature test is a bioassay for auxins.

Hint: Chapter plant growth and development, NCERT page no.247

**13.** (1)

Chapter: Evolution

Actually the wing of bat is homologous to flipper of whale. They have asked wings of birds. The most suitable answer is option 3 but it is a debatable question.

14. (4)

Chapter: Body fluids and Circulation

Blood flows under higher pressure in arteries than veins. So, blood pressure in Pulmonary artery will be higher than vena cava, but lower than aorta and carotid artery.

OUT of NCERT

15. (3)

Chapter: Human Reproduction

NCERT Page 51- 2nd para, 8th line

**16.** (2)

**Sol.:** Pachytene follows zygotene and is characterized by crossing over.

Hint: Chapter cell cycle cell division

NCERT page no.168

**17.** (3)

Hint: Chapter Biological classification

NCERT page no.20 and 21

18. (4)

Chapter: Movement and Locomotion

**OUT of NCERT** 

19. (3)

Sol.: Inhibin hormone is produced by granulosa cells of follicles in ovary and inhibit the secretion of FSH

Chapter: Human Reproduction

Inhibin is secreted by corpus luteum to inhibit FSH but not LH

20. (2)

**Sol.**: In Emphysema wall separating alveoli breaks and there is decrease in overall surface area available for gas exchange.

Chapter: Breathing and Exchange of Gases

NCERT Page 275, last para

21. (3)

**Sol.**: It follows X-linked / criss – cross inheritance.

Hint: Chapter Principles of Inheritance and Variation

NCERT page no.89

22. (4)

**Sol.:** *Sequoia* or redwood trees are among the tallest trees.

Hint: Chapter Plant kingdom

NCERT page no.38 and 39

23. (4)

Hint: Chapter Molecular Basis of Inheritance

NCERT page no.117

24. (2)

**Sol.:** Monohybrid mendelian cross

Hint: Chapter Principles of Inheritance

NCERT page no.73

**25.** (1)

**Sol.:** It is the root-knot nematode

Chapter: Biotechnology and Its applications

NCERT Page 210, 3rd para, 3rd Line

**26.** (1)

Sol.: Synapsis occurs during meiosis I between homologous chromosomes

Hint: Chapter Cell the unit of life

NCERT page no.137 and 138

**27.** (1)

Chapter: Human health and diseases

Cancer lines have large amount of telomerase. If mutation inhibits production of telomerase, the quantity of telomerase will be reduced.

**OUT of NCERT** 

28. (2)

**Sol.:** Chitin is a polymer of N-acetylglucosamine.

Hint: Chapter Biological classification

Not given in NCERT

29. (1)

**Sol.:** It is the rudimentary cotyledon.

Hint: Chapter Morphology of Flowering plants

NCERT page no.177

30. (2)

**Sol.:** Lichens are pioneer vegetation during xerarch.

Hint: Chapter Ecosystem

NCERT page no.250

31. (2)

Sol.: GnRH is secreted by hypothalamus & it acts on anterior pituitary to regulate production of FSH & LH

Chapter: Human Reproduction

Inhibin inhibits FSH from pituitary but doesn't inhibit GnRH from hypothalamus.

**OUT of NCERT** 

**32.** (1)

**Sol.:** Polio drops that are administered into the body contain attenuated pathogens.

Chapter: Human health and diseases

**OUT of NCERT** 

33. (3)

**Sol.:** Photosensitive compound in human retina consist of protein opsin & retinal

Chapter: Neural control and Co-ordination

NCERT Page 324- 4<sup>th</sup> Para, 4<sup>th</sup> Line

34. (3)

Hint: Chapter Anatomy of flowering plants

NCERT page no.89

**35.** (4)

**Sol.:** Parapodia are extensions of body wall in case of annelids

Chapter: Kingdom Animalia

Parapodia is a feature of Annelids

NCERT Page 53

**36.** (4)

**Sol.:** affinity of hemoglobin with oxygen decreases when pH decreases.

Chapter: Breathing and Exchange of Gases

High concentration of Hydrogen ions causes dissociation curve to shift towards right favouring breakdown of oxyhaemoglobin

NCERT Page 274

37. (2)

Chapter: Kingdom Animalia

Exception to option 1 is Prototherians (egg laying Mammals)

Exception to option 2 is Cyclostomes (jawless vertebrate)

Exception to option 3 is Crocodile (Reptile with 4 chambered heart)

38.

Hint: Chapter Principles of Inheritance and Variation

NCERT page no.75, 76, 77

39. **(3)** 

Sol.: Triglycerides consist of one glycerol and three fatty acid molecules.

Chapter: Biomolecules

A typical fat molecule or neutral fat or true fats or triglycerides consists of one glycerol and 3 fatty acid molecules.

NCERT Page 144- 2<sup>nd</sup> para

40. **(1)** 

Hint: Chapter Sexual Reproduction flowering plants

NCERT page no.21

41. **(1)** 

Glycine is simplest amino acid. Sulphur containing amino acids are cysteine & methionine. Sol.:

Chapter: Biomolecules

NCERT Page 145, 148

42. **(3)** 

Hint: Chapter Photosynthesis in higher plants

Not given in NCERT

43. **(2)** 

Hint: Chapter cell the unit of life, NCERT page no.129

44. **(3)** 

Hint: Chapter Ecosystem

Not given in NCERT

45.

Hint: Chapter Biodiversity and conservation, NCERT page no.264

46.

Due to development of proton gradient. Sol.:

Hint: Chapter Photosynthesis in higher plants

NCERT page no.214

47. **(3)** 

Hint: Chapter Molecular Basis of Inheritance

Not given in NCERT

48. **(4)** 

Hint: Chapter Biological classification

NCERT page no.19

49. (3)

**Sol.:** *Periplaneta americana* as spiral cleavage during embryonic development.

Chapter: Structural organization in Animals

N- acetyl glucosamine is chitin which forms exoskeleton in cockroach. Arthropods are metamerically segmented. Schizocoel is found in Annelids, Arthropods, and Molluscs. Cockroach shows indeterminate and spiral cleavage (also seen in Annelids and Molluscs)

**OUT of NCERT** 

50. (2)

**Sol.:** It is a system of rotating crops with legumes / grass pastures to improve soil structure and fertility.

Hint: Chapter Strategies for Enhancement in Food Production

Not given in NCERT

51. (1)

Hint: Chapter Microbes in human welfare, Not given in NCERT

52. (4)

**Sol.:** Urea is synthezied in liver from ammonia and carbon dioxide.

Chapter: Elimination of Nitrogenous waste

**OUT of NCERT** 

53. (1)

**Sol.:** It is an autosomal recessive disorder.

Hint: Chapter Principles of Inheritance and variation

NCERT page no.89 and 90

54. (1)

Sol.: Sphincter of Oddi guards the opening of hepatopancretic duct into the duodenum.

Chapter: Digestion and Absorption

NCERT Page 261, 2<sup>nd</sup> para, 3<sup>rd</sup> line

55. (3)

Hint: Chapter Cell the unit of life

NCERT page no.137 and 138

**56.** (1)

Hint: Sexual reproduction in flowering plant

NCERT page no.35

57. (2)

Hint: Chapter Morphology of Flowering Plants

NCERT page no. 81

**58.** (2)

Hint: Chapter Morphology of flowering plants

Not given in NCERT

59. (2)

**Sol.:** It is a thermostable enzyme.

Chapter: Biotechnology Principles and Processes

NCERT Page 203, 1st para, 8th Line

60. (4)

**Sol.:** Seen in Cactus, Opuntia etc.

Hint: Chapter Morphology of Flowering plants

Not given in NCERT

61. (4)

Chapter: Human health and diseases NCERT Page 153, 3<sup>rd</sup> para- 5<sup>th</sup> Line

**62. (2)** 

**Sol.:** It is written in Latin.

Hint: Chapter living world

NCERT page no.7

**63.** (1)

**Sol.:** The male gametes are motile / flagellated.

Hint: Chapter Plant Kingdom

NCERT page no.35 and 36

**64.** (1)

**Sol.:** Amniocentesis is not used for detection of cleft palate.

Chapter: Reproductive health

Cleft palate is a structural deformity which can be detected only on ultrasound.

OUT of NCERT

**65.** (3

**Sol.:** Parietal or oxyntic cells present in gastric glands of stomach secrete HCl.

Chapter: Digestion and Absorption

NCERT Page 262, 2<sup>nd</sup> para, 3<sup>rd</sup> line

**66. (3)** 

Hint: Chapter Cell cycle cell division

NCERT page no.165

67. (3)

Hint: Not given in NCERT

**68. (4)** 

Hint: Chapter Cell the unit of life

NCERT page no. 134

**69.** (1)

Hint: Chapter Biotechnology and its application

NCERT page no.211 Para 1, diagram 12.3

Insulin is a simple protein showing 3 chains A B and C connected by Disulphide bridges

- **70.** Incorrect Question (Bonus)
- 71. (1)
- **Sol.:** RNA is of low molecular weight.

Hint: Chapter Biological classification

NCERT page no.27

**72. (3)** 

Chapter: Evolution

NCERT Page 131, 1st para, 3rd line

73. (4)

**Sol.:** Level of LH and FSH increases gradually during the follicular phase.

Chapter: Human Reproduction

LH and FSH gradually increases during follicular phase.

NCERT Page 50, figure 3.9

74. (4)

Chapter: Kingdom Animalia

Birds are strictly oviparous

**75. (2)** 

Hint: Chapter Sexual Reproduction in Flowering Plants

NCERT page no.31

**76.** (1)

Hint: Chapter Sexual Reproduction in flowering plants

NCERT page no.38

77. (1)

**Sol.:** Sperm production continues after vasectomy

Chapter: Reproductive health

Vasectomy doesn't prevent spermatogenesis as it occurs in Testis.

NCERT Page 60, 5<sup>th</sup> Last line, 2<sup>nd</sup> para 1<sup>st</sup> line, Page 61- 2<sup>nd</sup> Para, 9<sup>th</sup>- 11<sup>th</sup> Line

**78.** (2)

Chapter: Chemical Control and Co-ordination

T3, T4 are derivatives of tyrosine. Estrogen, Progesterone, Cortisol and Cortisone are steroids.

OUT of NCERT

**79.** (1)

Hint: Chapter Environmental Issues

NCERT page no.275

80. (4)

Hint: Chapter Organisms and Population; NCERT page no.235

81. (3)

Chapter: Human Health and diseases

Mast cells are histaminic causing inflammation during asthma

NCERT Page 123, 2<sup>nd</sup> para

**82.** (4)

Hint: Chapter Morphology of Flowering plants

NCERT page no.74

83. (2)

Chapter: Biotechnology Principles and Process

NCERT Page 195, 5th para, 6th Line

84. (3)

Chapter: Organisms and Population; OUT of NCERT

**85.** (4)

Chapter: Evolution; OUT of NCERT

86. (3)

Hint: Chapter Principles of Inheritance and Variation

NCERT page no.90

87. (3)

Hint: Chapter Environmental Issues

NCERT page no.282

88. (4)

Hint: Chapter Environmental Issues

NCERT page no.285

89. (2)

Hint: Chapter Molecular Basis of Inheritance

NCERT page no.115

90. (3)


Hint: Chapter Ecosystem

Not given in NCERT

K-JEE

## **PHYSICS**

91.




Use C. of energy and centrifugal force at top most point.

92.







93.

93. (2) 
$$-\frac{GM}{r} = -5.4 \times 10^7$$

$$+\frac{GM}{r} = 6$$

(i) ÷ (ii)

$$\frac{\frac{GM}{r}}{\frac{GM}{r^2}} = r \cdot \frac{5.4 \times 10^6}{6}$$

 $r = 9 \times 10^3 \,\mathrm{km}$ 

94. **(1)** 


Total flux = n flux of each long

$$= 1000 \times 4 \times 10^{-3} = Li$$

$$L = \frac{10^3 \times 4 \times 10^{-3}}{4} = 1$$

95. **(2)** 

96. **(4)** 



$$T \sin \theta = \frac{kq^2}{x^2}$$

$$T \cos \theta = mg$$

$$\sin \theta = \frac{kq^2}{mgx^2} = \frac{x}{2l}$$

$$\Rightarrow q^2 = \frac{x^3 mg}{x2l}$$

$$q = \left(\frac{mg}{k\,2l}\right)^{\frac{1}{2}} x^{\frac{3}{2}}$$

$$\frac{dq}{dt} = \left(\frac{mg}{2 \times l}\right)^{\frac{1}{2}} \frac{3}{2} x^{\frac{1}{2}} \frac{dx}{dt}$$

$$\Rightarrow x^{1/2}$$
 v = constant

$$v \propto \chi$$
-1/2

# 97. (1)

Include energy =  $\frac{1}{2} \times 2 \times v^2$ 


Energy loss

$$\frac{1}{2} \times \left(\frac{2 \times 8}{2 + 8}\right) \times \left(v.0\right)^2$$

$$\frac{1}{2} \times \left(\frac{16}{10}\right) \times v^2$$

% loss = 
$$\frac{\frac{1}{2} \times \frac{16}{10} \times v^2}{\frac{1}{2} \times 2 \times v^2} \times 100 = \frac{1800}{2}$$

98. (4)



Particle is performing a circular motion.

$$I = I_0 - I_1$$

$$= \frac{MR^2}{2} - \left(\frac{M(R/2)^2}{4} + \frac{M(R/2)^2}{2} + \frac{M(R/2)^2}{4}\right)^2$$

$$= MR^2 \left[\frac{1}{2} - \frac{1}{32} - \frac{1}{16}\right]$$

$$= \frac{13MR^2}{32}$$

## 100.

$$= MR^{2} \left[ \frac{1}{2} - \frac{1}{32} - \frac{1}{16} \right]$$

$$= \frac{13MR^{2}}{32}$$
(3)
$$V_{e} = \sqrt{2gR}$$

$$= \sqrt{\frac{2GM}{R^{2}}R}$$

$$V_{e} = \sqrt{\frac{\frac{2GM}{4\pi}R^{2}}{3}}$$

$$V_{e} = \sqrt{2\left(\frac{4}{3}\right)\pi \times \rho R^{2}}$$

$$\frac{V_{e_{2}}}{V_{e_{1}}} = \sqrt{\left(\frac{\rho_{2}}{\rho_{1}}\right)\left(\frac{R_{2}}{R_{1}}\right)^{2}} = \sqrt{2 \times 2^{2}} = 2\sqrt{2}$$
(1)
$$\frac{\varepsilon_{1} + \varepsilon_{2}}{\varepsilon_{1} - \varepsilon_{2}} = \frac{5}{1} \quad \varepsilon_{1} + \varepsilon_{2} = 5\varepsilon_{1} - 5\varepsilon_{2}$$

$$6\varepsilon_{2} = 4\varepsilon_{1}$$

$$\frac{\varepsilon_{1}}{\varepsilon_{1}} = \frac{3}{2}$$

$$\frac{V_{e_2}}{V_{e_1}} = \sqrt{\left(\frac{\rho_2}{\rho_1}\right) \left(\frac{R_2}{R_1}\right)^2} = \sqrt{2 \times 2^2} = 2\sqrt{2}$$

## 101.

$$\frac{\varepsilon_1 + \varepsilon_2}{\varepsilon_1 - \varepsilon_2} = \frac{5}{1} \quad \varepsilon_1 + \varepsilon_2 = 5\varepsilon_1 - 5\varepsilon_2$$

$$6\varepsilon_2 = 4\varepsilon_1$$

$$\varepsilon_1 = 3$$

$$f = f_0 \left( \frac{V + V_0}{V - V_s} \right)$$

$$800 \left( \frac{330 + 0}{330 - 15} \right)$$

$$\frac{330}{315} \times 800$$

$$\frac{330}{315} \times 800 \left( \frac{330}{330} \right) = 838$$
Hz

103. **(1)** 

Using properties of OR and AND Gate

104.

 $\sin \theta = \frac{\lambda}{a}$  Position of first minima

$$\sin 30^\circ = \frac{5000}{a}$$

$$\sin\theta = \frac{3\lambda}{2a} = \frac{3 \times 5000}{2 \times 10,000}$$

 $\theta = \sin^{-1} \frac{3}{4}$  position of first secondary maxima.

105.

$$\frac{hc}{\lambda} = \frac{hc}{\lambda_0} + ev$$

$$\frac{hc}{2\lambda} = \frac{hc}{\lambda_0} + e\frac{v}{4}$$

$$\frac{\text{hc}}{2\lambda} = \frac{3\text{ev}}{4}$$

$$ev = \frac{2hc}{3\lambda}$$

$$\frac{hc}{\lambda} = \frac{hc}{\lambda} + \frac{2hc}{3\lambda}$$

$$\frac{hc}{\lambda_0} = \frac{hc}{3\lambda}$$

$$\lambda_0 = 3\lambda$$

106.

$$\frac{1}{2}mv^2 = \frac{kze^2}{r}$$

$$r \propto \frac{1}{m}$$

107.

$$(A) \rightarrow b, c;$$

$$(B) \rightarrow b, c$$

$$(C) \rightarrow b, d$$

$$(D) \rightarrow a, d$$

108. **(2)** 

$$\frac{1}{2} \times 10 \times 10^{-3} \times v^2 = 8 \times 10^{-4}$$


$$V^2 = 16 \times 10^{-2}$$

$$V = 4 \times 10^{-1} = 0.4 \,\text{m/s}$$

$$0.4 \times 0.4 = 0 + 2 \times a \times 4\pi \times \frac{6.4}{100}$$

$$a = 0.1 \text{ m/s}^2$$

109. **(3)** 



 $Cos\phi = 0$  so it will not consume energy

110.

Cos
$$\phi = 0$$
 so it will not consume energy

(3)
$$a_{c} = \frac{F}{M} \left[ \frac{1 + \frac{\Gamma}{R}}{1 + \frac{I_{c}}{MR^{2}}} \right] = g \sin \theta \left[ \frac{1}{1 + \frac{I_{c}}{MR^{2}}} \right]$$
Since sphere has less moment of inertia
So it reaches bottom first

(1)
$$l_{1}' = l_{1} \left( 1 + \alpha_{1} \Delta T \right)$$

$$l_{2}' = l_{2} \left( 1 + \alpha_{2} \Delta T \right)$$

$$l_{1}' = l_{1} + l_{1} \alpha_{1} \Delta T$$

$$l_{2}' = l_{2} + l_{2} \alpha_{2} \Delta T$$

$$l_{1}' - l_{2}' = l_{1} - l_{2} + \left( l_{1} \alpha_{1} - l_{2} \alpha_{2} \right) \Delta T$$

$$l_{1} \alpha_{1} = l_{2} \alpha_{2}$$
(1)

Since sphere has less moment of inertia So it reaches bottom first

111. **(1)** 

$$l_1' = l_1 \left( 1 + \alpha_1 \Delta T \right)$$

$$l_2' = l_2 \left( 1 + \alpha_2 \Delta T \right)$$

$$l_1' = l_1 + l_1 \alpha_1 \Delta T$$

$$l_2' = l_2 + l_2 \alpha_2 \Delta T$$

$$l_1' - l_2' = l_1 - l_2 + (l_1 \alpha_1 - l_2 \alpha_2) \Delta T$$

$$l_1 \alpha_1 = l_2 \alpha_2$$

112.

$$\frac{1}{V} - \frac{1}{-200} = \frac{1}{40}$$

$$\frac{1}{V} = \frac{1}{40} = \frac{1}{200} = \frac{5 - 1}{200} = \frac{4}{200} = \frac{1}{50}$$

$$V = 50 \text{ cm}$$

$$50 + 4 = 54$$
 cm

113. **(2)** 

$$\omega_f = \omega_i + \alpha t$$


$$= 0 + 2 \times 2 = 4$$

$$a_c = \omega^2 r = 4^2 \times \frac{1}{2} = 8$$

114. (4)

$$COP = \frac{T_2}{T_1 - T_2} = \frac{Heat \ extracted}{input \ work}$$

115. (3




116. (1)

Let intensity due to one slit is I

$$I_0 = 4 I$$

$$I = \frac{I_0}{4}$$




Phase diff. = 
$$\frac{2\pi}{\lambda} \cdot \frac{d}{D} \cdot \frac{d}{2} = \frac{2\pi}{\lambda} \cdot \frac{d}{10d} \cdot \frac{d}{2}d$$

$$\phi = \frac{2\pi}{\lambda} \cdot \frac{2\lambda}{10 \cdot 2} = \frac{\pi}{2}$$

$$I' = 4I\cos^2\frac{\phi}{2}$$

$$=4.\frac{I_0}{4}.\frac{1}{2}=\frac{I_0}{2}$$

# 117. (1)



$$ALdg = ApL \ n\rho g + Ah.\rho g$$

$$Ld = pL \ n\rho + h\rho$$

$$= pLn\rho + \rho L(1-p)$$

$$d = L\rho (np + 1 - p)$$

$$i = \frac{V}{R} = \frac{4 - (-6)}{1k\Omega} = \frac{10V}{1000\Omega} = 10^{-2} A$$

$$v = \sqrt{gR \frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}}$$

$$\frac{h_c}{\lambda} = \left[ \left( \frac{R}{h_c} \right) \left[ \frac{1}{4} - \frac{1}{\infty} \right] \right]$$

$$\frac{1}{\lambda} = \frac{R}{4} = \frac{10^7}{4}$$

Wave number =  $\frac{1}{\lambda}$ .  $0.25 \times 10^7$ 

$$V = At + Bt^2$$

$$V = At + Bt$$
$$= 0. t(A + Bt)$$

$$=0.t(A+bt)$$

$$t = -\frac{A}{B}$$

$$\frac{dx}{dt} = At + Bt^2$$

$$dx = \frac{At^2}{2} + \frac{Bt^3}{3} \bigg|_0^1$$

$$\frac{A}{2} + \frac{B}{3}$$


$$\frac{A \times 2t}{2} + \frac{B \times 2^{3}}{3}$$

$$2A + \frac{B8}{3}$$

$$2A - \frac{A}{2} + \frac{8B}{3} - \frac{B}{3}$$

$$\frac{3}{2}A + \frac{7B}{3}$$

**123. (3)** 



$$\mu = \delta + A = i_1 + e$$

$$\delta+60^\circ=90^\circ$$

$$\delta = 30^{\circ}$$

$$G = \frac{\sin\left(\frac{A+\delta_m}{2}\right)}{\sin\left(\frac{A}{2}\right)} = \frac{\sin 45^{\circ}}{\sin 30^{\circ}} = \frac{1/\sqrt{2}}{1/2} = \sqrt{2}$$


124.


$$v_{rms} \propto \sqrt{T}$$

$$\frac{v}{200} = \sqrt{\frac{400}{300}}$$

$$v = \frac{200 \times 2}{\sqrt{3}} = \frac{400}{\sqrt{3}}$$

125. **(4)** 





$$(2\lambda + 1)\frac{\lambda}{2} = \ell$$

$$\lambda = \frac{\mu}{2n+1}f = c$$

$$f = \frac{(2n+1)}{2\ell}$$

$$\ell = \frac{(26+1)c}{2f}$$

$$N = 0$$

and 
$$N = 1$$

127. (2) 
$$\lambda_e = \frac{h}{\sqrt{2mE}}$$

$$\lambda_p = \frac{hc}{E}$$

$$\frac{\lambda_e}{\lambda_p} = c \left(\frac{E}{2m}\right)^{1/2}$$

$$P = \vec{F} \cdot \vec{v}$$

$$= (2t\hat{i} + 2t^2\hat{j})N$$

$$= 2t^3\hat{i} + 3t^5\hat{j}$$

$$\vec{a} = (2t\hat{i} + 3t^2\hat{j}) \text{ m/s}^2$$

$$\vec{v} = t^2\hat{i} + t^3\hat{j}$$

$$Q = 0 \Rightarrow t^{20}, t = \frac{a}{2B}$$

$$dH = i^{2}Rdt$$

$$i = \frac{ds}{dt} = a - 2b - f$$

$$H = \int dH \int_{0}^{2b} (a - 2bt)^{2} Rdt$$

$$= \left[ \frac{(a - 2bt)^{3}}{-6b} \right]_{0}^{b/2}$$

$$= \frac{-1}{6b} \left[ 0 - a^{3} \right] R$$

$$= \frac{a^{3}R}{6b}$$

$$v_{\text{gain}} = \beta \frac{R_L}{R_{BE}}$$

Power gain =  $\beta^2 \frac{R_L}{R_{BE}}$ 


$$\frac{1}{2} \times mgh = mL$$

$$F = F_{AB} - F_{LD}$$

$$= \frac{\mu_0 I}{2\pi \frac{L}{2}} iL + \frac{\mu_0 I}{2\pi \frac{3L}{2}} iL$$

$$= \frac{\mu_0 Ii}{2\pi} \left(2 - \frac{2}{3}\right)$$

$$=\frac{2\mu_0 Ii}{3\pi}$$



$$\frac{\lambda_2}{\lambda_1}$$

$$v \propto \sqrt{7}$$

$$\frac{v_A}{v_B} = \sqrt{\frac{m_2}{m_2 + m_1}} \frac{m_1 + m_2}{m_2}$$

**134.** (4

$$\lambda_m T = b$$

and 
$$U \propto T^4$$

$$T_1 > T_2 > T_3$$

$$U_1 > U_2 > U_3$$

135. (1)

Theory

## **CHEMISTRY**

- 136. (3
- **Sol.** The physical and chemical adsorption is accompanied with decrease in FREE ENERGY, ENTHALPY and ENTROPY.
- 137. (2
- **Sol.** Hydrogen ion concentration in pure water at  $298 \text{ K} = 10^{-7} \text{m}$ . Reduction potential of hydrogen electrode is given by.

$$E_{H^{+}/H_{2}} = -\frac{0.0591}{2} log \frac{P_{H_{2}}}{\left[H^{+}\right]^{2}}$$

$$E_{H^+/H_2} = 0$$
 if  $P_{H_2} = [H^+]^2 = 10^{-14}$ atm

- 138. (1)
- **Sol.** The addition of a catalyst during a chemical reaction alters the activation energy.
- 139. (3)
- **Sol.** Reactions (a), (b) and (c) are elimination, substitution and addition respectively.
- 140. (2)

**Sol.** Aldehyde reacts with a primary amine to form schiff base

$$R - CH = O + H_2N - R' \longrightarrow R - CH = N - R' + H_2O$$

- 141. (2)
- **Sol.** Arylamines are less basic than alkylamines because the lone pair of electrons on N-atom is involved in resonance with the benzene ring.
- 142. (2)
- Sol. Let the initial moles of each of  $H_2$  and  $O_2$  be 1. Number of moles of  $H_2$  diffused in certain time = 0.5. Number of moles of  $O_2$  diffused (say x) in the same time is given by  $\frac{r_{O_2}}{r_{H_3}} = \frac{x}{0.5} = \sqrt{\frac{2}{32}} = \frac{1}{4} \Rightarrow x = \frac{1}{8}$
- 143. (1)
- **Sol.** The staggered conformation of ethane is more stable than eclipsed conformation because staggered conformation has no torsional strain.
- **144.** (3 & 4)
- Sol. The correct order of first ionisation enthalpy is B < C < O < NThe correct order of electron gain enthalpy is I < Br < F < CI
- 145. (2)
- **Sol.** Rate of a first order reaction at 10 min and 20 min is given by

$$R_{10} = k[A]_{10} = 0.04 \text{ mol } L^{-1} \text{ s}^{-1}$$

$$R_{20} = k[A]_{20} = 0.03 \text{ mol } L^{-1}s^{-1}$$

$$\frac{\left[A\right]_{10}}{\left[A\right]_{20}} = \frac{4}{3} = e^{(20-10)k} = e^{10k}$$

On solving, 
$$k = \frac{2.303 \times 0.125}{10} s^{-1}$$

Half life, 
$$f_{1/2} = \frac{0.693 \times 10}{2.303 \times 0.125} = 24.1s$$

- 146. (2)
- Sol. Copper reacts with conc. HNO<sub>3</sub> to give  $Cu(NO_3)_2$  and  $NO_2$  $Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + 2H_2O$
- 147. (4
- **Sol.** In a protein molecule various amino acids are linked together by peptide bond.
- 148. (2)
- **Sol.** Fog is a colloid of liquid dispersed in gas.
- 149. (2)
- Sol. (a) cyanide process is applicable for Au

- (b) Froth floatation process is applicable for sulphide ores like ZnS.
- (c) Electrolytic reduction is done for 'Al'
- (d) Zone refining process is used for the extraction of ultrapure metals like Ge, Si, etc.
- **150.** (1)
- **Sol.** Sucrose is a non reducing sugar.
- 151. (3)
- **Sol.** The sugar in RNA is ribose and the sugar in DNA is 2'-deoxyribose.
- 152. (4)
- **Sol.**  $\Delta G = \Delta H T\Delta S$

For a spontaneous process,  $\Delta G = -ve$ . So,  $\Delta H < 0$  and  $\Delta S > 0$ 

- 153. (2
- **Sol.** Phosphinic acid (H<sub>3</sub>PO<sub>2</sub>) is monoprotic acid and phosphonic acid (H<sub>3</sub>PO<sub>3</sub>) is diprotic acid.
- 154. (3)
- **Sol.** If s and s' are the solubilities of MY and NY<sub>3</sub> respectively, then

$$s = \sqrt{6.2 \times 10^{-13}} = 7.87 \times 10^{-7} \,\mathrm{M}$$

$$s' = \left(\frac{6.2 \times 10^{-13}}{27}\right)^{1/4} = 3.89 \times 10^{-4} \,\mathrm{M}$$

- 155. (2
- **Sol.** Novalgin is analgesic, rest are antibiotics.
- **156.** (1)

$$CH_3 - C \equiv C^{\circ}$$

**Sol.** Hybridization sp

The lone pair is present in 'sp' hybridized orbital of 'C'.

- 157. (3)
- **Sol.** Stability of conjugate base :

$$ClO_4^- > ClO_3^- > ClO_2^- > ClO_3^-$$

- **158.** (1)
- **Sol.**  $K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \longrightarrow Cr_2(SO_4)_3 + K_2SO_4 + H_2O_4$
- 159. (2)
- **Sol.** According to VSEPR theory: lone pair lone pair bond pair bond pair bond pair bond pair
- **160.** (1)
- **Sol.**  $m_s = \pm \frac{1}{2}$  for 2 electrons having rest same quantum numbers.

According to Pauli's exclusion principle, no two electrons of same spin can occupy the same orbital.

- **161.** (Bonus)
- **Sol.**  $CaC_2 + N_2 \longrightarrow CaCN_2 + C$

- 162. **(2)**
- Sol. Natural rubber is polymer of Cis-isoprene units.
- 163.
- Bond dissociation energy Sol.

$$Cl_2 > Br_2 > F_2 > I_2$$

- 164. **(2)**
- $ROH \xrightarrow{NaH} RO^{-} \xrightarrow{R'-X} R O R'$ Sol.

The given reaction is Williamson ether synthesis.

165. **(3)** 

**Sol.** 
$$d = \frac{Z \times M}{a^3 \times N_A}$$

$$a^{3} = \frac{2 \times 6.94}{530 \times 10^{-3} \times 6.02 \times 10^{23}} = \frac{13.88}{530 \times 10^{-3} \times 6.02 \times 10^{23}}$$

$$a = 352 \text{pm}$$
(2)
$$\frac{\Gamma^{+}}{\Gamma^{-}} = 0.54 \implies \text{C.N} = 6$$
(2)
$$\frac{P^{\circ} - P_{s}}{P_{s}} = \frac{n_{\text{solute}}}{n_{\text{solvent}}} = \frac{6.5 / \text{M}}{100 / 18}$$

$$\frac{760 - 732}{732} = \frac{6.5 \times 18}{100 \times \text{M}}$$

$$M = \frac{6.5 \times 18 \times 732}{28 \times 100} = 30.58$$

$$\Delta T_{b} = K_{b}.m = 0.52 \times \frac{6.5}{30.58} = 1.1^{\circ}\text{C}$$
(1)
$$T_{b} = 101.1^{\circ}\text{C}$$

166. **(2)** 

**Sol.** 
$$\frac{r^+}{r^-} = 0.54 \implies C.N = 6$$

**167.** 

Sol. 
$$\frac{P^{\circ} - P_{s}}{P_{s}} = \frac{n_{\text{solute}}}{n_{\text{solvent}}} = \frac{6.5 / M}{100 / 18}$$

$$\frac{760 - 732}{732} = \frac{6.5 \times 18}{100 \times M}$$

$$M = \frac{6.5 \times 18 \times 732}{28 \times 100} = 30.58$$

$$\Delta T_b = K_b.m = 0.52 \times \frac{6.5}{30.58} = 1.1^{\circ} \text{C}$$

$$T_b = 101.1^{\circ} C$$

- 168. **(1)**
- Sol. Stability of half filled 'f' sub-shell.
- 169. (1 & 2)
- **170. (2)**

**Sol.** 
$$H - C \equiv CH \xrightarrow{(1)NaNH_2/liq.NH_3} H - C \equiv C \xrightarrow{(2)CH_3CH_2Br} H - C \equiv C - CH_2 - CH_3$$

171. **(1)** 

**Sol.** 
$$\frac{d \ln P}{dT} = \frac{\Delta H_{v}}{RT^{2}}$$
 [Clausius – Clapeyron equation]

Sol. 
$$y_B = \frac{p_B}{p_T} = \frac{p_B^0 X_B}{p_A^0 X_A + p_B^0 X_B}$$


$$y_{\text{Benzene}} = \frac{12.8 \times \frac{1}{2}}{\frac{1}{2} (12.8 + 3.85)}$$

$$y_{\text{Benzene}} = 0.77$$

$$y_{\text{toluene}} = 0.23$$

$$\therefore y_{\text{Benzene}} > y_{\text{Toluene}}$$

173. (3)



It has no plane of sym and centre of sym.

**Sol.** Cis-cyclopenta-1, 2 diol forms a ring like structure called isopropylidene derivative with acetone in acidic medium, while trans-cyclopenta-1, 2 diol can't form ring like structure.

$$\begin{array}{c}
OH \\
OH
\end{array}$$

$$\begin{array}{c}
O \\
CH_3 - C - CH_3 / H^{\dagger}
\end{array}$$

$$\begin{array}{c}
O \\
CH
\end{array}$$

175. (1

Carbonyl compounds with  $\alpha$ -hydrogen atom readily equilibrates into its enol form due to acidic nature of  $\alpha$ -hydrogen atom. This is known as keto-enol tautomerism.

176. (3)

Sol. Molecule  $CH_4$   $NH_3$   $H_2O$  Bond angle  $109.5^{\circ}$   $107^{\circ}$   $104.5^{\circ}$ 

177. (2)

**Sol.** (a)  $XeF_6: H = sp^3d^3$ ; 6B.P + 1 L.P - distorted octahedral

(b)  $XeO_3 : H = sp^3 ; 3B.P + 1 LP - Pyramidal$ 

(c)  $XeOF_4$ :  $H = sp^3d^2$ ; 5B.P + 1 LP - square pyramidal

(d)  $XeF_4$ :  $H = sp^3d^2$ : 4B.P + 2LP - square planar

178. (3)

- Sol. Addition of large amount of KHSO<sub>4</sub> to the nitrating mixture reduces the rate of nitration by lowering the conc. of  $NO_2^+$  ion.
- 179. (4)
- **Sol.** Ca<sup>2+</sup> ions are important in maintaining the regular heart beat.
- 180. (4)
- Sol. Since Fe in  $[Fe(CO_4)^{2-}]$  has 2 –ve charges, its tendency to use its filled orbital to overlap with the vacant anti-bonding MO of CO is high, so C O bond length in this complex is the longest.

